

KNOTS AND CLASSICAL 3-GEOMETRIES

Tze-Chuen Toh* and Malcolm R. Anderson†

**Department of Theoretical Physics
Research School of Physical Sciences and Engineering
The Australian National University
Canberra ACT 0200 AUSTRALIA*

*and
†Department of Mathematics
Edith Cowan University
Mt. Lawley WA 6050 AUSTRALIA*

ABSTRACT. It has been conjectured by Rovelli that there is a correspondence between the space of link classes of a Riemannian 3-manifold and the space of 3-geometries (on the same manifold). An exact statement of his conjecture will be established and then verified for the case when the 3-manifold is compact, orientable and closed.

PACS. 04.60, 02.40.

Mathematics Subject Classifications (1991). 53C80, 54P47 + 35.

(This article will appear in the January, 1995 edition of the Journal of Mathematical Physics.)

1. Introduction.

In [6, p. 1661], Rovelli sketched a proof showing how a certain collection of n -loops, which he called *weaves*, are related to the flat 3-metric. He then conjectured that perhaps there exists a relationship between n -loops,¹ for $n < \infty$, and 3-metrics. The relationship between n -loops, for $n < \infty$, and 3-metrics will not be answered in this paper (and so, it still remains an open question); however, what will be shown in this article is that there exists a precise relationship between 3-geometries and a subset of $\aleph_0$ -knots, where an n -knot is defined to be an equivalence class of n -loops

*e-mail: tct105@rsphy2.anu.edu.au

†e-mail: m.anderson@cowan.edu.au

¹An n -loop $\gamma \stackrel{\text{def}}{=} \{\gamma^1, \dots, \gamma^n\}$ is just a subset of the loop space consisting of n loops; i.e., γ^i , for each $i = 1, \dots, n$, are (distinct) closed curves in Σ .

under (smooth) ambient isotopies (cf. §3). The approach given here is entirely different to that outlined by Rovelli in [6]: tersely, Rovelli introduced a lattice spacing on the 3-manifold—the distance between parallel non-intersecting curves which defines a weave in 3-space—to obtain his conclusion regarding weaves and flat metrics; this, in turn, motivated his conjecture. Here, no such assumptions will be made and the results are purely ‘topological’.

The attention here will be focused on compact, Riemannian 3-manifolds. In this paper, the term *Riemannian metric* means a smooth, non-degenerate, symmetric, covariant 2-tensor that is *positive-definite* on the base manifold. The fact that the 3-manifold is separable is crucial in the construction: this, at least, explains why $\aleph_0$ -loops are used rather than n -loops for $n < \infty$. The main interest in Rovelli’s conjecture is that it will provide a tentative physical interpretation of the loop representation of quantum gravity [7]: it yields a possible insight into the interweaving of topology and geometry at the quantum level. More will be said in section 5.

In all that follows, the spatial (Riemannian) 3-manifold, denoted by Σ , is assumed to be smooth, orientable, closed and compact; $\mathbb{R}_+ \stackrel{\text{def}}{=} \{s \in \mathbb{R} \mid s \geq 0\}$ and $I \stackrel{\text{def}}{=} [0, 1]$. Lastly, let $\text{Diff}^+(\Sigma)$ denote the group of smooth, orientation-preserving, diffeomorphisms on Σ . An overview of this paper runs as follows: section 2 introduces the required notations and definitions, whilst in section 3, the property of the space of $\aleph_0$ -knots of a subset of $\aleph_0$ -loops will be examined. This space will establish the sought for correspondence between topology and geometry. In section 4, a variant of the Rovelli conjecture will be formulated precisely and then verified; then, in section 5, some interesting speculations regarding the results established in §4 will be outlined.

2. Preliminary Definitions and Notations.

Let $\Omega_\Sigma = \{\gamma : I \rightarrow \Sigma \mid \gamma(0) = \gamma(1), \gamma \text{ continuous}\}$ be the space of loops in Σ and equip it with the compact-open topology. Fix a Riemannian metric $\hat{q}$ on Σ and let $\hat{d} : \Sigma \times \Sigma \rightarrow \mathbb{R}_+$ be the distance function on Σ induced by $\hat{q}$. Then, the metric $d_\Omega : \Omega_\Sigma \times \Omega_\Sigma \rightarrow \mathbb{R}_+$ defined by

$$d_\Omega(\gamma, \eta) \stackrel{\text{def}}{=} \sup_{t \in I} \hat{d}(\gamma(t), \eta(t))$$

induces a metric topology on Ω_Σ which is compatible with its compact-open topology [1, p. 263, theorem 4.2.17].

2.1. Remark. Since for each (admissible) Riemannian metric q on Σ ,² the d_q -topology—where d_q is the distance function induced on Σ by q —coincides with the manifold topology, it follows that all the metrics d_q are equivalent to one another. Hence, all the metrics d_Ω are also equivalent.

Let $\tilde{\mathcal{L}}_\Sigma \subset \Omega_\Sigma$ be the space of piecewise smooth loops in Σ endowed with the subspace topology. Next, quotient away the constant loops—i.e., $\gamma(I) = \{x_\gamma\}$, some $x_\gamma \in \Sigma$ —in $\tilde{\mathcal{L}}_\Sigma$ as follows. Define an equivalence relation $\tilde{\mathcal{R}} \subset \tilde{\mathcal{L}}_\Sigma \times \tilde{\mathcal{L}}_\Sigma$ such that $\forall (\gamma, \eta) \in \tilde{\mathcal{R}}$, γ and η are constant loops in $\tilde{\mathcal{L}}_\Sigma$. Let $\mathcal{L}_\Sigma \stackrel{\text{def}}{=} \tilde{\mathcal{L}}_\Sigma / \tilde{\mathcal{R}}$ be the quotient space and $\tilde{\pi} : \tilde{\mathcal{L}}_\Sigma \rightarrow \mathcal{L}_\Sigma$ the natural map. It is clear that $\tilde{\mathcal{R}}$ is closed in $\tilde{\mathcal{L}}_\Sigma \times \tilde{\mathcal{L}}_\Sigma$. Furthermore, observe from the construction that if $\mathcal{L}_0 = \{\gamma \in \tilde{\mathcal{L}}_\Sigma \mid \gamma \text{ is a constant loop}\}$, then $\tilde{\pi}|(\tilde{\mathcal{L}}_\Sigma - \mathcal{L}_0)$ coincides with the inclusion map $\tilde{\mathcal{L}}_\Sigma - \mathcal{L}_0 \hookrightarrow \tilde{\mathcal{L}}_\Sigma$; that is, $\tilde{\pi}|(\tilde{\mathcal{L}}_\Sigma - \mathcal{L}_0) = \text{id}_{\tilde{\mathcal{L}}_\Sigma}|(\tilde{\mathcal{L}}_\Sigma - \mathcal{L}_0)$.

2.2. Lemma. $\mathcal{L}_0$ is closed and nowhere dense in $\tilde{\mathcal{L}}_\Sigma$.

Proof. Let $\{\gamma_n\}_n$ be a sequence in $\mathcal{L}_0$ which converges to $\gamma_0 \in \tilde{\mathcal{L}}_\Sigma$. By definition, $\gamma_n(I) = x_n \in \Sigma \forall n$ and Σ compact Hausdorff imply that $\exists x_0 \in \Sigma$ and a subsequence $\{x_{n_k}\}_k \subset \{x_n\}_n$ such that $x_{n_k} \rightarrow x_0$. Since $\forall \varepsilon > 0, \exists N_\varepsilon > 0$ such that $d_\Omega(\gamma_n, \gamma_0) = \sup_{t \in I} \hat{d}(\gamma_n(t), \gamma_0(t)) = \sup_{t \in I} \hat{d}(x_n, \gamma_0(t)) < \varepsilon \forall n > N_\varepsilon$, it follows at once that $\gamma_0(t) \equiv x_0$ on I and $\mathcal{L}_0$ is thus closed, where $d_\Omega|_{\tilde{\mathcal{L}}_\Sigma \times \tilde{\mathcal{L}}_\Sigma}$ is denoted by d_Ω for simplicity.

Finally, to complete the proof, suppose that the interior $\mathcal{L}_0^\circ \neq \emptyset$. Then, for any fixed $\eta \in \mathcal{L}_0^\circ$, there exists a neighbourhood $N_\eta = \bigcap_{i=1}^n M(K_i, O_i)$, for some $n < \infty$ —where $M(K_i, O_i) \stackrel{\text{def}}{=} \{\gamma \in \tilde{\mathcal{L}}_\Sigma \mid K_i \subset I \text{ is compact, } \gamma(K_i) \subset O_i, O_i \subset \Sigma \text{ is open}\}$ —such that $N_\eta \subset \mathcal{L}_0^\circ$. However, $\eta \in N_\eta$ and $\eta(I) = x_\eta$ for some $x_\eta \in \Sigma \Rightarrow O \equiv \bigcap_{i=1}^n O_i \neq \emptyset$. Let $L_\eta = \{\gamma \in \tilde{\mathcal{L}}_\Sigma \mid \gamma(I) \subset O\} - \mathcal{L}_0$. Evidently, $L_\eta \neq \emptyset$, and in particular, $L_\eta \cap N_\eta \neq \emptyset$, which is a contradiction. Hence, $\mathcal{L}_0^\circ \equiv \emptyset$, as required. $\square$

The neighbourhood base of $0_\Sigma \equiv \tilde{\pi}(\gamma) \forall \gamma \in \mathcal{L}_0$ can now be constructed. It follows from the quotient topology and lemma 2.2 that for each neighbourhood N_{0_Σ} of 0_Σ , $\tilde{\pi}^{-1}(N_{0_\Sigma})$ must be a neighbourhood of $\mathcal{L}_0$. Hence, from the definition of $\tilde{\pi}$, the neighbourhood base of 0_Σ consists precisely of subsets $\tilde{\pi}(N)$, where N is a neighbourhood of $\mathcal{L}_0$ in $\tilde{\mathcal{L}}_\Sigma$. Explicitly, a neighbourhood of 0_Σ is of the form

²The topology on Σ and its differentiable structure are of course assumed fixed throughout the discussion.

$\bigcup_{\eta \in \mathcal{L}_0} \tilde{\pi}(N_\eta)$, where N_η is a neighbourhood of η in $\tilde{\mathcal{L}}_\Sigma$. Notice however, that $\tilde{\pi}$ is not an open map. For given any neighbourhood N_η of $\eta \in \mathcal{L}_0$, $\tilde{\pi}^{-1} \circ \tilde{\pi}(N_\eta) = N_\eta \cup \mathcal{L}_0$ which is neither closed nor open (by lemma 2.2) if $\mathcal{L}_0 \not\subset N_\eta$. Nevertheless, for each neighbourhood N of $\mathcal{L}_0$, $\tilde{\pi}(N)$ is a neighbourhood of 0_Σ .

2.3. Lemma. $\tilde{\pi} : \tilde{\mathcal{L}}_\Sigma \rightarrow \mathcal{L}_\Sigma$ is closed.

Proof. To establish this claim, it is enough to show that $\tilde{\pi}$ maps closed neighbourhoods C_η of $\eta \in \mathcal{L}_0$ into closed neighbourhoods of 0_Σ . Invoking the quotient topology, it will suffice to verify that $\tilde{\pi}^{-1} \circ \tilde{\pi}(C_\eta)$ is closed in $\tilde{\mathcal{L}}_\Sigma$. Since $\tilde{\pi}^{-1} \circ \tilde{\pi}(C_\eta) = C_\eta \cup \mathcal{L}_0$ by definition, lemma 2.2 yields the desired result. $\square$

2.4. Theorem. $\mathcal{L}_\Sigma$ is metrizable.

Proof. By corollary 2.3, it will suffice to show that $\mathcal{L}_\Sigma$ is first countable [1, p. 285, theorem 4.4.17], and from the definition of $\tilde{\pi}$, it is enough to verify that 0_Σ has a countable neighbourhood base, since each $\gamma \in \mathcal{L}_\Sigma - \{0_\Sigma\}$ has a countable neighbourhood base (by definition). Let $\mathfrak{B}_\eta = \{B_{\frac{1}{n}}(\eta) \mid n \in \mathbb{N}\}$ be a countable neighbourhood base of $\eta \in \mathcal{L}_0$ in $\tilde{\mathcal{L}}_\Sigma$. Since, by definition, each subset of the form $\bigcup_{\eta \in \mathcal{L}_0} \tilde{\pi}(B_{\frac{1}{n}}(\eta))$ defines a neighbourhood of 0_Σ , it follows that the collection $\mathfrak{B}_{0_\Sigma}$ defined by

$$\mathfrak{B}_{0_\Sigma} = \left\{ \bigcup_{\eta \in \mathcal{L}_0} \tilde{\pi}(B_{\frac{1}{n}}(\eta)) \mid n \in \mathbb{N} \right\}$$

is a countable neighbourhood base of 0_Σ . Hence, $\mathcal{L}_\Sigma$ is metrizable, as claimed. $\square$

In the following account, call a curve in Σ a *q-geodesic* if it is a geodesic in Σ relative to the Riemannian metric q . Also, if γ, η are curves such that $\gamma(1) = \eta(0)$, then define $\gamma * \eta$ by

$$\gamma * \eta(t) = \begin{cases} \gamma(2t) & \text{for } 0 \leq t \leq \frac{1}{2}, \\ \eta(2t - 1) & \text{for } \frac{1}{2} \leq t \leq 1. \end{cases}$$

2.5. Definition. Let $\gamma \in \mathcal{L}_\Sigma$. Then, γ is said to be a *piecewise geodesic loop* if there exists a Riemannian metric q on Σ and n smooth q -geodesics $\gamma_1, \dots, \gamma_n : I \rightarrow \Sigma$, $1 \leq n < \infty$, such that $\gamma = \gamma_1 * \dots * \gamma_n$.³

Let Γ_2^+ denote the space of (smooth) Riemannian metrics on Σ (endowed with the compact C^∞ -topology⁴) and $D_\Sigma \subset \Sigma$ a countably dense subset of Σ . Now,

³Note that each γ_i in γ is still a q -geodesic with respect to its new parametrization $[\frac{i-1}{n}, \frac{i}{n}]$, as the reparametrization $\gamma_i(t) \rightarrow \gamma_i(nt - i + 1) \equiv \gamma[\frac{i-1}{n}, \frac{i}{n}]$ is clearly an affine transformation.

⁴The compact C^∞ -topology is defined in the appendix.

define $\mathcal{M}_\infty[q]$, for each $q \in \Gamma_2^+$, to be the set of all countably infinite multi-loops $\gamma = \{\gamma^i \mid i \in \mathbb{N}\}$ satisfying the following two properties:

- (1) for each i , $\gamma^i \in \mathcal{L}_\Sigma$ is a piecewise (affinely parametrized) q -geodesic loop in Σ ,
- (2) the subset γ is in bijective correspondence with D_Σ under the map $\gamma^i \mapsto \gamma^i(0)$.

Finally, set $\mathcal{M}_\infty[\Gamma_2^+] = \bigcup_{q \in \Gamma_2^+} \mathcal{M}_\infty[q]$. An immediate consequence of the definition is the following two observations. Suppose $\gamma \in \mathcal{M}_\infty[q] \cap \mathcal{M}_\infty[q']$. Let $\Gamma(q)$ and $\Gamma(q')$ be the Riemannian connections of q and q' respectively. Fix an admissible atlas $\{(U_\alpha, \psi_\alpha)\}_\alpha$ on Σ . Then, with respect to each chart U_α ,

$$(\ddot{\gamma}_\alpha^i)^\ell + \Gamma_\alpha(q)_{kj}^\ell (\dot{\gamma}_\alpha^i)^k (\dot{\gamma}_\alpha^i)^j \stackrel{\text{a.e.}}{=} 0 \quad \text{and} \quad (\ddot{\gamma}_\alpha^i)^\ell + \Gamma_\alpha(q')_{kj}^\ell (\dot{\gamma}_\alpha^i)^k (\dot{\gamma}_\alpha^i)^j \stackrel{\text{a.e.}}{=} 0$$

on $\gamma^i(I) \cap U_\alpha$ for each i (no summation over α , obviously), where $F(t) \stackrel{\text{a.e.}}{=} 0$ means $F(t) = 0$ on I apart from a *finite* number of points in I . Hence, $(\Gamma_\alpha(q)_{kj}^\ell - \Gamma_\alpha(q')_{kj}^\ell)(\dot{\gamma}_\alpha^i)^k (\dot{\gamma}_\alpha^i)^j \stackrel{\text{a.e.}}{=} 0 \quad \forall \gamma^i \in \gamma$ and α . Thus, by property (2), $\Gamma(q)_{kj}^\ell(x) \equiv \Gamma(q')_{kj}^\ell(x)$ on a dense subset of Σ as $\overline{\bigcup\{\gamma^i(I) \mid \gamma^i \in \gamma\}} \equiv \Sigma$ by (2). Hence, invoking the continuity of $\Gamma(h)$ for $h = q, q'$, it follows at once that $\Gamma(q) \equiv \Gamma(q')$ on Σ . Now, with respect to local coordinate basis, $\Gamma(q)_{kj}^i = \frac{1}{2}q^{ih}(\partial_k q_{hj} + \partial_j q_{hk} - \partial_h q_{kj})$ (and likewise for q'); consequently, q and q' are related homothetically; that is, $\exists c > 0$ constant such that $q' = cq$.⁵ More generally, q, q' are related by some coordinate transformation, as is shown below.

Let $f : \Sigma \rightarrow \Sigma$ be a smooth diffeomorphism, where $\Sigma = (\Sigma, q)$ and set $\Sigma_f = f(\Sigma) \stackrel{\text{def}}{=} (\Sigma, (f^{-1})^*q)$. Clearly, if $\gamma : I \rightarrow \Sigma$ is a q -geodesic, then $\gamma : I \rightarrow \Sigma_f$ is an $(f^{-1})^*q$ -geodesic in Σ_f and conversely, by symmetry (as isometries map geodesics into geodesics). Hence, in view of these two observations, each $\gamma \in \mathcal{M}_\infty[\Gamma_2^+]$ is assigned to a unique 3-geometry of Σ , where the space of *3-geometries* is defined to be the quotient space $\mathcal{Q} = \Gamma_2^+ / \text{Diff}^+(\Sigma)$. Recall that each element $[q] \in \mathcal{Q}$ is defined by $[q] = \{f^*q \mid f \in \text{Diff}^+(\Sigma)\}$. Let $\pi_+ : \Gamma_2^+ \rightarrow \mathcal{Q}$ denote the natural projection. Then, π_+ is open [2, p. 317, §3.1] and $\mathcal{Q}$ is a second countable, metrizable space [2, p. 318, theorem 1].

As a converse remark, notice that if Σ were not separable or that $\gamma_q = \{\gamma_q^i \mid i \in \mathbb{N}\}$ were not chosen to satisfy (2), γ_q need not uniquely determine $[q] \in \mathcal{Q}$. For want of a better term, call $\mathcal{M}_\infty[\Gamma_2^+]$ the space of *piecewise geodesic* $\aleph_0$ -loops.

⁵Note trivially that as q, q' are positive-definite, $c < 0$ is not an admissible solution.

Now, a suitable topology can be defined on this space. To do this, let L_∞ be the set of affinely parametrized, piecewise geodesic loops in Σ and let L_Σ^∞ denote the countably infinite set-theoretic product of L_Σ . Define an equivalence relation $R_\Sigma \subset L_\Sigma^\infty \times L_\Sigma^\infty$ by

$$R_\Sigma \stackrel{\text{def}}{=} \{(\gamma, \gamma') \in L_\Sigma^\infty \times L_\Sigma^\infty : [\gamma] = [\gamma']\},$$

where $[\eta] \stackrel{\text{def}}{=} \{\eta^i \mid i \in \mathbb{N}\}$ is just the set of components of $\eta \stackrel{\text{def}}{=} (\eta^i)_{i=1}^\infty$. Let $\pi_\Sigma : L_\Sigma^\infty \rightarrow M_\Sigma \stackrel{\text{def}}{=} L_\Sigma^\infty / R_\Sigma$ denote the natural map. Then clearly, as a subset, $\mathcal{M}_\infty[\Gamma_2^+] \subset M_\Sigma$.

Now, let $M_\infty \subset L_\Sigma^\infty$ be a subset satisfying

- (i) for each $\gamma \stackrel{\text{def}}{=} (\gamma^i)_{i=1}^\infty$, $\gamma^i \neq \gamma^j \ \forall i \neq j$,
- (ii) $\pi_\Sigma(M_\infty) = \mathcal{M}_\infty[\Gamma_2^+]$.

It is clear from the definition of M_∞ that there exists a family of subsets $M_\sigma \subset M_\infty$ such that

- (a) $M_\infty = \bigcup_\sigma M_\sigma$,
- (b) $M_\sigma \cap M_{\sigma'} = \emptyset \ \forall \sigma \neq \sigma'$,
- (c) $\pi_\Sigma|_{M_\sigma} : M_\sigma \rightarrow \mathcal{M}_\infty[\Gamma_2^+]$ is a (set-theoretic) bijection.

Let $h_\sigma \stackrel{\text{def}}{=} \pi_\Sigma|_{M_\sigma}$ and for each $\gamma \in \mathcal{M}_\infty[\Gamma_2^+]$, set $\gamma_\sigma = h_\sigma^{-1}(\gamma) \in M_\sigma$.⁶ The subsets M_σ admit suitable metrics to be constructed below. Firstly, fix a finite atlas $\mathfrak{A}$ on Σ and let $\tilde{d}_\Omega$ be a metric on $\mathcal{L}_\Sigma$ compatible with its quotient topology. Then, for any pair $\gamma, \eta \in M_\sigma$, let $d_\sigma(\gamma, \eta) \stackrel{\text{def}}{=} \sup_i \tilde{d}_\Omega(\gamma^i, \eta^i) + \sup_i \tilde{d}'_\Omega(\gamma^i, \eta^i)$, where

$$\tilde{d}'_\Omega(\gamma^i, \eta^i) \stackrel{\text{def}}{=} \text{ess sup} \{ \|D^k \gamma^i(t) - D^k \eta^i(t)\| : t \in I, k \geq 1 \}$$

with sup running over all relevant (finite) charts $(U, \varphi) \in \mathfrak{A}$, ess denoting that the expression $\|D^k \gamma^i(t) - D^k \eta^i(t)\|$ is *not* defined only on a *finite* (possibly zero) set of points in I wherein γ^i and η^i are not differentiable, and $D^k \gamma^i(t)$ denotes the k th differential of γ^i at t in abused notation. It is routine to verify that d_σ is indeed a metric. In all that follows, M_σ will be endowed with the d_σ -topology.

2.6. Remark. Let $\overline{\mathfrak{A}}$ denote the maximal atlas of Σ and define a topology on M_σ to be generated by subbasic open sets $N_\varepsilon(\gamma; (U_{\alpha(i)}, \varphi_{\alpha(i)})_{i=1}^\infty, K)$ in M_σ to be constructed below, where $K \subset I$ is compact, $\gamma^i(K) \subset U_{\alpha(i)}$ and $(U_{\alpha(i)}, \varphi_{\alpha(i)}) \in$

⁶The subscript σ on γ_σ will be omitted should no confusion arise from the context.

$\overline{\mathfrak{A}} \forall i$. Denote $\{\alpha(i) \mid 1 \leq i \leq \infty\}$ by α and $(U_{\alpha(i)}, \varphi_{\alpha(i)})_i$ by $(U, \varphi)_\alpha$ for notational simplicity, and let

$$\tilde{d}'_{\sigma\alpha K}(\gamma^i, \eta^i) \stackrel{\text{def}}{=} \{\|D^k \varphi_{\alpha(i)} \circ \gamma^i(t) - D^k \varphi_{\alpha(i)} \circ \eta^i(t)\| : t \in K, k \geq 1\}$$

whenever $\gamma^i(K), \eta^i(K) \subset U_{\alpha(i)} \forall i$. Then, for a fixed $\gamma \in M_\sigma$ such that $\gamma^i(K) \subset U_{\alpha(i)} \forall i$, let $N_\varepsilon(\gamma; (U, \varphi)_\alpha, K) \stackrel{\text{def}}{=} \{\eta \in M_\sigma \mid \tilde{d}_{\sigma\alpha K}(\gamma, \eta) < \varepsilon, \eta^i(K) \subset U_{\alpha(i)} \forall i\}$, where

$$\tilde{d}_{\sigma\alpha K}(\gamma, \eta) \stackrel{\text{def}}{=} \sup_i \tilde{d}_\Omega(\gamma^i, \eta^i) + \sup_i \tilde{d}'_{\sigma\alpha K}(\gamma^i, \eta^i).$$

It can be shown that this topology is equivalent to the $\tilde{d}_\sigma$ -topology on M_σ . In particular, the $\tilde{d}_\sigma$ -topologies on M_σ defined relative to any two finite atlases of Σ are equivalent. Hence, in this sense, the $\tilde{d}_\sigma$ -topology is well-defined as it does not depend on the choice of *finite* atlas $\mathfrak{A}$ on Σ .

A topology on $\mathcal{M}_\infty[\Gamma_2^+]$ can now be constructed. Firstly, notice that the spaces M_σ and $M_{\sigma'}$ are homeomorphic for each pair σ, σ' —define $h_{\sigma\sigma'} : M_\sigma \rightarrow M_{\sigma'}$ by $\gamma_\sigma \mapsto \gamma_{\sigma'}$, where $h_\sigma(\gamma_\sigma) = \gamma = h_{\sigma'}(\gamma_{\sigma'})$. The existence of $h_{\sigma\sigma'}$ follows immediately from conditions (i) and (c). Hence, it is possible to endow $\mathcal{M}_\infty[\Gamma_2^+]$ with a topology such that each $h_\sigma : M_\sigma \rightarrow \mathcal{M}_\infty[\Gamma_2^+]$ defines a homeomorphism. This will be the topology imposed on $\mathcal{M}_\infty[\Gamma_2^+]$. As an aside, if M_∞ is given the sum topology, $M_\infty \stackrel{\text{def}}{=} \bigoplus_\sigma M_\sigma$, then $h : M_\infty \rightarrow \mathcal{M}_\infty[\Gamma_2^+]$ given by $h|_{M_\sigma} \stackrel{\text{def}}{=} h_\sigma$ defines a continuous open surjection.

3. The Space of $\aleph_0$ -Knots of $\mathcal{M}_\infty[\Gamma_2^+]$.

First of all, some notations and elementary properties of the space of equivalence $\aleph_0$ -loop classes will be established. Let $\mathcal{G}_a^+$ be the set of (smooth) orientation-preserving, ambient isotopies on Σ . That is, $\mathcal{G}_a^+ \subset C^\infty(\Sigma \times I, \Sigma \times I)$ is the following set:

$$\{F : \Sigma \times I \rightarrow \Sigma \times I \mid F(x, t) \stackrel{\text{def}}{=} (F_t(x), t), F_0 = \text{id}_\Sigma, F_t \in \text{Diff}^+(\Sigma) \forall t \in I\}$$

and define composition $\circ$ on $\mathcal{G}_a^+$ by

$$(F' \circ F) : (x, t) \mapsto (F'_t \circ F_t(x), t).$$

Then, clearly, $F' \circ F \in \mathcal{G}_a^+$ and $1_{\Sigma \times I} \stackrel{\text{def}}{=} \text{id}_\Sigma \times \text{id}_I \in \mathcal{G}_a^+$. It is straight forward to check that $\langle \mathcal{G}_a^+, \circ \rangle$ forms a group under $\circ$, where the inverse F^{-1} of $F = (F_t, \text{id}_I)$

is defined to be (F_t^{-1}, id_I) . In particular, $\circ$ is compatible with the compact C^∞ -topology on $\mathcal{G}_a^+$ —cf. [3, p. 64, ex. 9]. Moreover, since $\text{Diff}^+(\Sigma)$ is closed in the group $\text{Diff}(\Sigma)$ of smooth diffeomorphisms endowed with the compact C^∞ -topology (as it is a subgroup of $\text{Diff}(\Sigma)$), $\mathcal{G}_a^+$ is also closed in $C^\infty(\Sigma \times I, \Sigma \times I)$ (with respect to the compact C^∞ -topology).

If $\gamma, \eta \in \mathcal{L}_\Sigma$ are any pair of loops and γ is ambiently isotopic to η under some $F \in \mathcal{G}_a^+$, denote this by $F : \gamma \simeq \eta$. Now, given any pair of $\aleph_0$ -loops $\gamma, \eta \in \mathcal{M}_\infty[\Gamma_2^+]$, define an equivalence relation R generated by $\simeq$ on $\mathcal{M}_\infty[\Gamma_2^+]$ as follows:

$$\gamma \simeq \eta \iff \exists F \in \mathcal{G}_a^+ \text{ such that } F \cdot \gamma = \eta,$$

where $F \cdot \gamma \stackrel{\text{def}}{=} \{F_1 \circ \gamma^1, F_1 \circ \gamma^2, \dots\}$ and $F : \gamma^i \simeq \eta^i \forall i$. Then, the space $\mathcal{K}[\Gamma_2^+]$ of equivalence classes of $\aleph_0$ -loops in $\mathcal{M}_\infty[\Gamma_2^+]$ is defined to be the quotient space $\mathcal{M}_\infty[\Gamma_2^+]/\mathcal{G}_a^+$. Henceforth, for simplicity, the term (*piecewise geodesic*) $\aleph_0$ -*knot* will mean an element of the quotient space $\mathcal{K}[\Gamma_2^+]$; that is, an $\aleph_0$ -knot denotes an equivalence class of $\aleph_0$ -loops under a smooth, orientation-preserving, ambient isotopy. The space $\mathcal{K}[\Gamma_2^+]$ will be called the $(\aleph_0, \Gamma_2^+)$ -*knot space* of $\mathcal{M}_\infty[\Gamma_2^+]$. Let $\kappa_\infty : \mathcal{M}_\infty[\Gamma_2^+] \rightarrow \mathcal{K}[\Gamma_2^+]$ denote the natural map, where $\mathcal{K}[\Gamma_2^+]$ is endowed with the quotient topology.

3.1. Lemma. *The natural projection $\kappa_\infty : \mathcal{M}_\infty[\Gamma_2^+] \rightarrow \mathcal{K}[\Gamma_2^+]$ is open.*

Proof. A sketch of the proof will be given. To see that κ_∞ is an open mapping, it is enough to note that for each open subset $N \subset \mathcal{M}_\infty[\Gamma_2^+]$,

$$\kappa_\infty^{-1} \circ \kappa_\infty(N) = \bigcup_{F \in \mathcal{G}_a^+} F \cdot N,$$

where $F \cdot N = \{F \cdot \gamma \mid \gamma \in N\}$. Since $F \cdot N$ is open in $\mathcal{M}_\infty[\Gamma_2^+]$, as F defines a homeomorphism from $\mathcal{M}_\infty[\Gamma_2^+]$ onto itself, the quotient topology implies that $\kappa_\infty^{-1} \circ \kappa_\infty(N)$, and hence κ_∞ , must also be open. $\square$

3.2. Proposition. *$\mathcal{K}[\Gamma_2^+]$ is Hausdorff.*

Proof. By lemma 3.1, it will suffice to show that the equivalence relation R generated by $\simeq$ is closed in $\mathcal{M}_\infty[\Gamma_2^+] \times \mathcal{M}_\infty[\Gamma_2^+]$ [4, p. 98, theorem 11]. Let $\{(\gamma_n, \eta_n)\}_n$ be a sequence in R which converges in $\mathcal{M}_\infty[\Gamma_2^+] \times \mathcal{M}_\infty[\Gamma_2^+]$ to (γ_0, η_0) . By definition, $\exists$ a sequence $\{F_n\}_n$ in $\mathcal{G}_a^+$ such that $F_n : \gamma_n \simeq \eta_n$ for each n . So, $(\gamma_n, F_n \cdot \gamma_n) \rightarrow (\gamma_0, \eta_0) \Rightarrow F_n \cdot \gamma_n \rightarrow \eta_0$ and $\gamma_n \rightarrow \gamma_0$, and hence implying that

$\{F_n\}_n$ is a convergent sequence in $\mathcal{G}_a^+$. Consequently, $\mathcal{G}_a^+$ is closed implies that $F_n \rightarrow F_0 \in \mathcal{G}_a^+$ for some F_0 . Whence, $\eta_0 \equiv F_0 \cdot \gamma_0$ and R is thus closed, as desired. $\square$

In the interest of simplicity, call $\gamma \in \mathcal{M}_\infty[\Gamma_2^+]$ a *piecewise $(\aleph_0, q)$ -geodesic loop* whenever the 3-metric q is required to be specified.

3.3. Lemma. *Let $\gamma, \tilde{\gamma} \in \mathcal{M}_\infty[\Gamma_2^+]$ be piecewise $(\aleph_0, q)$ - and $(\aleph_0, \tilde{q})$ -geodesic loops respectively. If $\gamma \simeq \tilde{\gamma}$, then $\exists f \in \text{Diff}^+(\Sigma)$ such that $q = f^*\tilde{q}$.*

Proof. Let $F \in \mathcal{G}_a^+$ be an ambient isotopy of γ and $\tilde{\gamma}: F \cdot \gamma = \tilde{\gamma}$. Then, evidently, $\tilde{\gamma}$ is a piecewise $(\aleph_0, (F_1^{-1})^*q)$ -geodesic. However, $\tilde{\gamma}$ is also a piecewise $(\aleph_0, \tilde{q})$ -geodesic; hence, by §2 (2), $\exists f \in \text{Diff}^+(\Sigma)$ such that $\tilde{q} = f^*q$, as required. $\square$

4. $\aleph_0$ -Knots and Classical Geometry.

In this section, the relationship between the equivalence classes of $\aleph_0$ -loops in Σ and the (classical) geometries admissible on Σ will be studied. This correspondence can be easily sought simply by noting that each element in $\mathcal{M}_\infty[\Gamma_2^+]$ corresponds to a unique 3-geometry $[q]$ of Σ by construction. The modified form of Rovelli's Conjecture can now be formulated.

4.1. Theorem. *There exists a continuous, open surjection $\hat{\chi} : \mathcal{M}_\infty[\Gamma_2^+] \rightarrow \mathcal{Q}$ given by $\gamma_q \mapsto [q]$, where γ_q is a (piecewise) $(\aleph_0, q)$ -geodesic loop and $q \in [q]$.*

Proof (Sketch). Firstly, $\hat{\chi}$ is well-defined from the definition of $\mathcal{M}_\infty[\Gamma_2^+]$. Secondly, the surjective property of $\hat{\chi}$ is also clear. Thirdly, in this proof, Γ_2^+ will be identified with its image under the (topological) imbedding $j^\infty : \Gamma_2^+ \hookrightarrow C(\Sigma, J^\infty[p_\Sigma])$.⁷ So, $\mathcal{Q} \equiv j^\infty \Gamma_2^+ / \text{Diff}^+(\Sigma)$ and $\pi_+ : j^\infty \Gamma_2^+ \rightarrow \mathcal{Q}$.

Now, fix some $\gamma_0 \in \mathcal{M}_\infty[\Gamma_2^+]$ and let $N(q_0) = \bigcap_{i=1}^n M(K_i, (\pi_\Sigma^{n_i})^{-1}(U^{n_i}))$ be a neighbourhood of q_0 in Γ_2^+ , where $n_i \in \mathbb{N}$, $n < \infty$ and $q_0 \in \hat{\chi}(\gamma_0) = [q_0]$ is a representative of the q_0 -equivalence class. Set $N([q_0]) = \pi_+(N(q_0))$. Then, $\tilde{N}([q_0]) \stackrel{\text{def}}{=} \pi_+^{-1}(N([q_0])) = \bigcup \{ f^* \circ N(q_0) \mid f \in \text{Diff}^+(\Sigma) \}$, where $f^* \circ N(q_0) \stackrel{\text{def}}{=} \{ f^*q \mid q \in N(q_0) \}$. Let $D_\varepsilon(\gamma_0)$ be an ε -neighbourhood of γ_0 defined by $B_\varepsilon(h_\sigma^{-1}(\gamma_0)) = h_\sigma^{-1}(D_\varepsilon(\gamma_0)) \forall \sigma$. Then, $\forall \eta \in D_\varepsilon(\gamma_0)$, $\tilde{d}_\Omega(\gamma_{0\sigma}^i, \eta_\sigma^i) + \tilde{d}'_\Omega(\gamma_{0\sigma}^i, \eta_\sigma^i) < \varepsilon \forall i$ and σ , where $h_\sigma^{-1}(\gamma) \stackrel{\text{def}}{=} \gamma_\sigma$.

⁷The notations used here—the C^∞ -jets and compact C^∞ -topology—can be found in the appendix.

Next, observe from the definition that

$$(*) \quad (\ddot{\gamma}^i)^\ell + \Gamma(q)_{kj}^\ell (\dot{\gamma}^i)^k (\dot{\gamma}^i)^j \stackrel{\text{a.e.}}{=} 0 \quad \forall i \in \mathbb{N} \text{ and } \ell = 1, 2, 3,$$

where $\Gamma(q)$ is a Riemannian connection determined by the 3-metric q (with the connection coefficients written with respect to the natural frame for simplicity). So, by choosing $\varepsilon > 0$ to be sufficiently small, and by fixing *any* σ —and setting $\gamma_0^i = \gamma_0^{\sigma(i)}$, $\eta^i = \eta^{\sigma(i)}$ —it follows that $|\ddot{\eta}^i - \ddot{\gamma}_0^i| < \varepsilon$ and $|\dot{\eta}^i - \dot{\gamma}_0^i| < \varepsilon$ (almost everywhere), and in particular, using $(*)$,

$$\begin{aligned} & \left| (\ddot{\gamma}_0^i)^\ell + \Gamma(q_\eta)_{kj}^\ell (\dot{\gamma}_0^i)^k (\dot{\gamma}_0^i)^j \right| \\ & \stackrel{\text{a.e.}}{=} \left| (\ddot{\eta}^i + \mathcal{O}(\varepsilon))^\ell + \Gamma(q_\eta)_{kj}^\ell (\dot{\eta}^i + \mathcal{O}(\varepsilon))^k (\dot{\eta}^i + \mathcal{O}(\varepsilon))^j \right| \\ & \stackrel{\text{a.e.}}{=} \left| (\ddot{\eta}^i)^\ell + \Gamma(q_\eta)_{kj}^\ell (\dot{\eta}^i)^k (\dot{\eta}^i)^j + \mathcal{O}(\varepsilon) \right| \\ & \sim \mathcal{O}(\varepsilon) \text{ a.e. on } I, \end{aligned}$$

where $q_\eta \in \hat{\chi}(\eta)$. Whence, $|(\ddot{\gamma}_0^i)^\ell + \Gamma(q_\eta)_{kj}^\ell (\dot{\gamma}_0^i)^k (\dot{\gamma}_0^i)^j| \stackrel{\text{a.e.}}{=} |(\ddot{\gamma}_0^i)^\ell + \Gamma(q_0)_{kj}^\ell (\dot{\gamma}_0^i)^k (\dot{\gamma}_0^i)^j - (\ddot{\gamma}_0^i)^\ell - \Gamma(q_\eta)_{kj}^\ell (\dot{\gamma}_0^i)^k (\dot{\gamma}_0^i)^j| = |(\Gamma(q_0) - \Gamma(q_\eta))_{kj}^\ell (\dot{\gamma}_0^i)^k (\dot{\gamma}_0^i)^j| \sim \mathcal{O}(\varepsilon)$ a.e. (from above) $\forall i \in \mathbb{N} \Rightarrow |\Gamma(q_\eta)_{kj}^\ell - \Gamma(q_0)_{kj}^\ell|$ is small on Σ for each fixed ℓ, k, j whenever $\varepsilon > 0$ is small enough by appealing to §2 (2) and the continuity of Γ . Thus, from $\Gamma(q)_{kj}^\ell \stackrel{\text{def}}{=} \frac{1}{2} q^{\ell h} (\partial_k q_{hj} + \partial_j q_{hk} - \partial_h q_{kj})$ (in the natural frame), it follows that $\exists f \in \text{Diff}^+(\Sigma)$ such that $f^* q_\eta$ and q_0 , together with their k th derivatives, must be close to one another: $f^* q_\eta(K_i) \subset (\pi_\Sigma^{n_i})^{-1}(U^{n_i}) \quad \forall i = 1, \dots, n$. So, $f^* q_\eta$ and hence q_η must both belong to $\tilde{N}([q_0])$ for $\varepsilon > 0$ sufficiently small. Whence, $\hat{\chi}(D_\varepsilon(\gamma_0)) \subset N([q_0])$, and the continuity of $\hat{\chi}$ follows.

Finally, to conclude this proof, observe that for any $\gamma \in \mathcal{M}_\infty[\Gamma_2^+]$, $\hat{\chi}^{-1} \circ \hat{\chi}(\gamma) = \{f \circ \gamma \mid f \in \text{Diff}^+(\Sigma)\}$, where $f \circ \gamma \stackrel{\text{def}}{=} \{f \circ \gamma^1, f \circ \gamma^2, \dots\}$. Hence, for any ε -neighbourhood $D_\varepsilon(\gamma)$,

$$\hat{\chi}^{-1} \circ \hat{\chi}(D_\varepsilon(\gamma)) = \bigcup_{f \in \text{Diff}^+(\Sigma)} f \circ D_\varepsilon(\gamma),$$

and $\hat{\chi}$ is thus open, as desired. $\square$

In spite of the divergent approach given here with Rovelli's original idea, the following corollary could perhaps be christened as the *weak Rovelli conjecture* inasmuch as the notion of relating knots with geometry originated from Rovelli [6].

4.2. Corollary (Weak Rovelli Conjecture). *The map $\hat{\chi}$ induces a continuous, open surjection $\chi : \mathcal{K}[\Gamma_2^+] \rightarrow \mathcal{Q}$ given by $[\gamma_q] \mapsto \hat{\chi}(\gamma_q)$, where $\gamma_q \in \kappa_\infty^{-1}([\gamma_q])$ is any fixed representative.*

Proof. This map χ is well-defined by lemma 3.3. The result now follows immediately from theorem 4.1, lemma 3.1 and the commutativity of the following diagram:

$$\begin{array}{ccc} \mathcal{M}_\infty[\Gamma_2^+] & \xrightarrow{\hat{\chi}} & \mathcal{Q} \\ \kappa_\infty \downarrow & & \downarrow \text{id} \\ \mathcal{K}[\Gamma_2^+] & \xrightarrow{\chi} & \mathcal{Q}. \end{array}$$

□

Two comments regarding theorem 4.1 and its corollary are now in order. Firstly, it is certainly evident that if Σ be separable (which, here, it is in any case!), then it is sufficient to characterized its 3-geometries by the $\aleph_0$ -loops in $\mathcal{M}_\infty[\Gamma_2^+]$ since, by construction, $\overline{\{\gamma^i(0) \mid i \in \mathbb{N}\}} \equiv \Sigma$, whereas n -loops, for $n < \infty$ (using this construction), are not sufficient to determine the 3-geometry uniquely (as might well be expected): cf. §2.5 for a detailed account.

Secondly, it has been established elsewhere—cf. for example, [7, p. 132, §5.1] using the diffeomorphism constraints of general relativity (in the loop representation)—that functionals on $\mathcal{L}_\Sigma$ which describe gravitational states are constant on the $\mathcal{G}_a^+$ -orbits of $\mathcal{L}_\Sigma$: $\psi[\gamma] = \psi[\gamma'] \ \forall \gamma, \gamma' \in [\gamma]$, where $\psi : \mathcal{L}_\Sigma \rightarrow \mathbb{C}$ is a loop functional. However, surprisingly, this condition follows immediately from corollary 4.2. This can be easily seen as follows. Functionals on Γ_2^+ that describe gravitational states are those which are invariant under $\text{Diff}^+(\Sigma)$: i.e., they are essentially functionals on $\mathcal{Q}$. Let $C(\mathcal{Q}, \mathbb{C})$ be the set of continuous functionals on $\mathcal{Q}$ and let $C(\mathcal{K}[\Gamma_2^+], \mathbb{C})$ be the set of functionals on $\mathcal{K}[\Gamma_2^+]$. Then, $\forall \tilde{\Psi} \in C(\mathcal{Q}, \mathbb{C})$, $\tilde{\Psi} \circ \chi \in C(\mathcal{K}[\Gamma_2^+], \mathbb{C})$; that is, $\chi^*(C(\mathcal{Q}, \mathbb{C})) \subset C(\mathcal{K}[\Gamma_2^+], \mathbb{C})$, and the assertion thus follows.

This concludes the classical description of $\aleph_0$ -knots and their relationship with 3-geometries.

5. Discussion.

In this final section, a possible physical interpretation—albeit a highly speculative one!—regarding knots and gravity will be sketched. As was pointed out before,

the separability of Σ guarantees that $\hat{\chi}$ in theorem 4.1 remains well-defined. Furthermore, as classically, gravity—or equivalently, the 4-metric—of space-time is determined by the distribution of matter in the universe via Einstein’s field equations, gravity is a ‘global’ concept. In this sense, if n -loops can describe gravity in any way, then, provided that space-time be separable, loops that will best describe it are $\aleph_0$ -loops. Indeed, a judicious choice of $\aleph_0$ -loops—such as those given in the preceding sections—enables one to recover the underlying Riemannian 3-manifold Σ simply because $\overline{\{\gamma_q^i(0) \mid i \in \mathbb{N}\}} = \Sigma$, and $\hat{\chi}(\gamma_q) = [q]$. In the light of this observation, it is not unreasonable to conclude that gravity is the result of the way 3-space (and hence, space-time) is knotted, where (Σ, q) is said to be $[\gamma]$ -*knotted* if $\chi([\gamma]) = [q]$. And since χ is not one-one, Σ can be knotted in two $\mathcal{G}_a^+$ -inequivalent ways and yet give rise to the same gravitational configuration (determined by χ). In short, having determined $\mathcal{M}_\infty[\Gamma_2^+]$ from Σ , each element in $\mathcal{M}_\infty[\Gamma_2^+]$ contains the necessarily information to reconstruct Σ .

To conclude with a speculative note on the quantum aspect of a knot $[\gamma]$, one might heuristically interpret a *knot state* $|\gamma\rangle$ to correspond to the pair $[(\Sigma, q)]$, where $[(\Sigma, q)] \stackrel{\text{def}}{=} \{(\Sigma, q) \mid q \in \chi([\gamma])\}$. In particular, $|\gamma\rangle$ is associated with a particular 3-geometry $\chi([\gamma])$. Thus, $|\gamma\rangle$ corresponds to the global degrees of freedom of gravity: and since gravitons are associated with the local degrees of freedom of gravity, it has *no direct* relationship with a knot state. In the full quantum theory, it is quite reasonable to expect that $|\gamma\rangle$ will not span a Hilbert space due to the highly non-linear nature of gravity and the violation of the asymptotic completeness condition. Hence, a knot state most probably cannot be interpreted in the usual quantum field theoretic sense in that it lies in some Hilbert space, although it is tempting to conjecture that the knot states lie in some $\aleph_0$ -dimensional smooth Kähler manifold.

Acknowledgment. The first author thanks S. Scott, L. Tassie and P. Leviton for some fruitful conversations.

APPENDIX

A. Compact C^∞ -Topology.

The definition of a *compact C^∞ -topology* will be reviewed [5, pp. 32–33, §§4.1–4.3]. Let $J^n[\Sigma]$ be the space of C^n -jets from Σ into Σ and denote an element

in $J^n[\Sigma]$ by either $j^n(f(x))$ or $[f, x]_n$ (which ever proves more convenient). Fix an atlas $\mathfrak{A}_\Sigma = \{(U_\alpha, \psi_\alpha)\}_{\alpha \in \Lambda}$ on Σ and set $\mathfrak{A}_\Sigma(U_\alpha) = \{U \subset U_\alpha \mid U \text{ open}\}$. Then, $\mathfrak{B}_\Sigma = \bigcup_\alpha \mathfrak{A}_\Sigma(U_\alpha)$ forms a base for Σ . Let $J^0[\Sigma] = \Sigma \times \Sigma$ and let $\pi_1^0 : J^1[\Sigma] \rightarrow J^0[\Sigma]$ by $j^1\phi(p) \mapsto (p, \phi(p))$. Set $U_{\alpha\alpha'}^1 \equiv (\pi_1^0)^{-1}(U_\alpha \times U_{\alpha'})$ and define $p_\pm^1 : J^1[\Sigma] \rightarrow \Sigma$ by $p_+^1 : j^1\phi(p) \mapsto \phi(p)$ and $p_-^1 : j^1\phi(p) \mapsto p$. Then, it is clear that $U_{\alpha\alpha'}^1 = (p_-^1)^{-1}(U_\alpha) \cap (p_+^1)^{-1}(U_{\alpha'})$. Finally, let $\mathfrak{A}_{\alpha\alpha'}^1 = \{(\pi_1^0)^{-1}(U \times U') \mid U \times U' \subset U_\alpha \times U_{\alpha'} \text{ open}\}$. Then, $\mathfrak{B}^1 = \bigcup_{\alpha, \alpha'} \mathfrak{A}_{\alpha\alpha'}^1$ forms a base for $J^1[\Sigma]$. Following [8, p. 94, definition 4.1.5], define $\Psi_{\alpha\alpha'}^1 : U_{\alpha\alpha'}^1 \cong {}^3B_{\varepsilon_\alpha}(x_\alpha) \times {}^3B_{\varepsilon_{\alpha'}}(x_{\alpha'}) \times {}^{N_1}B_{\varepsilon_1}(x_1)$ by

$$[\phi, p]_1 \mapsto (\psi_\alpha(p), \psi_{\alpha'}(\phi(p)), D_\alpha(j^1\phi(p))),$$

where ${}^nB_\varepsilon(x)$ is an open ε -ball in $\mathbb{R}^n$ and $D_\alpha j^1\phi(p) \stackrel{\text{def}}{=} \left\{ \frac{\partial}{\partial x_\alpha^{i_1}} \phi_{\alpha\alpha'}(\psi_\alpha(p)) \right\}_{i_1}$ for some $N_1 \in \mathbb{N}$ such that $D_{\alpha\alpha'} : U_{\alpha\alpha'}^1 \cong {}^{N_1}B_{\varepsilon_1}(x_1)$ and $\phi_{\alpha\alpha'} \stackrel{\text{def}}{=} \psi_{\alpha'} \circ \phi \circ \psi_\alpha^{-1}$. The pair $(U_{\alpha\alpha'}^1, \Psi_{\alpha\alpha'}^1)$ defines a chart on $J^1[\Sigma]$. Denote $\Psi_{\alpha\alpha'}^1$ symbolically by $\psi_\alpha \times \psi_{\alpha'} \times D_\alpha$.

Now, define $p_\pm^2 : J^2[\Sigma] \rightarrow \Sigma$ by $p_-^2 : j^2\phi(p) \mapsto p$ and $p_+^2 : j^2\phi(p) \mapsto \phi(p)$. Furthermore, define $\pi_2^0 : J^2[\Sigma] \rightarrow J^0[\Sigma]$ by $\pi_2^0(j^2\phi(p)) = (p, \phi(p))$ and let $U_{\alpha\alpha'}^2 \stackrel{\text{def}}{=} (\pi_2^0)^{-1}(U_\alpha \times U_{\alpha'})$. Then, $U_{\alpha\alpha'}^2 \equiv (p_-^2)^{-1}(U_\alpha) \cap (p_+^2)^{-1}(U_{\alpha'})$. And as with the case for $J^1[\Sigma]$, the pair $(U_{\alpha\alpha'}^2, \Psi_{\alpha\alpha'}^2)$ defines a chart in $J^2[\Sigma]$, where $\Psi_{\alpha\alpha'}^2 \stackrel{\text{def}}{=} \psi_\alpha \times \psi_{\alpha'} \times D_\alpha \times D_\alpha^2$ and $D_\alpha^2 : U_{\alpha\alpha'}^2 \cong {}^{N_2}B_{\varepsilon_2}(x_2)$, for some $\varepsilon_2 > 0$ and some $N_2 \in \mathbb{N}$, is defined by $D_\alpha^2(j^2\phi(p)) \stackrel{\text{def}}{=} \left\{ \frac{\partial^2}{\partial x_\alpha^{i_1} \partial x_\alpha^{i_2}} \phi_{\alpha\alpha'}(\psi_\alpha(p)) \right\}_{i_1 \leq i_2}$. Also, define $\pi_2^1 : J^2[\Sigma] \rightarrow J^1[\Sigma]$ by $[\phi, p]_2 \mapsto [\phi, p]_1$. Then, by definition, $\pi_2^0 = \pi_1^0 \circ \pi_2^1$ and $\pi_2^1(U_{\alpha\alpha'}^2) = U_{\alpha\alpha'}^1$.⁸ Finally, let $\mathfrak{A}_{\alpha\alpha'}^2 = \{(\pi_2^0)^{-1}(U \times U') \mid U \times U' \subset U_\alpha \times U_{\alpha'} \text{ open}\}$; then, $\mathfrak{B}^2 = \bigcup_{\alpha, \alpha'} \mathfrak{A}_{\alpha\alpha'}^2$ forms a base for $J^2[\Sigma]$.

By induction, given $J^n[\Sigma]$, $(\pi_n^0)^{-1}(U_\alpha \times U_{\alpha'}) = (p_-^n)^{-1}(U_\alpha) \cap (p_+^n)^{-1}(U_{\alpha'})$ and $\pi_n^{n-1}(U_{\alpha\alpha'}^n) = U_{\alpha\alpha'}^{n-1}$. Furthermore, the pair $(U_{\alpha\alpha'}^n, \Psi_{\alpha\alpha'}^n)$ forms a chart on $J^n[\Sigma]$ as follows: $\Psi_{\alpha\alpha'}^n \stackrel{\text{def}}{=} \psi_\alpha \times \psi_{\alpha'} \times \prod_{i=1}^n D_\alpha^i$, where

$$D_\alpha^\ell : [\phi, p]_\ell \mapsto \left\{ \frac{\partial^\ell \phi_{\alpha\alpha'} \circ \psi_\alpha(p)}{\partial x_\alpha^{i_1} \dots \partial x_\alpha^{i_\ell}} \right\}_{i_1 \leq \dots \leq i_\ell} \in \mathbb{R}^{N_\ell}$$

with some $N_\ell \in \mathbb{N}$ such that $D_\alpha^\ell(U_{\alpha\alpha'}^\ell) = {}^{N_\ell}B_{\varepsilon_\ell}(x_\ell)$. Tersely, $\Psi_{\alpha\alpha'}^n : U_{\alpha\alpha'}^n \cong {}^3B_{\varepsilon_\alpha}(x_\alpha) \times {}^3B_{\varepsilon_{\alpha'}}(x_{\alpha'}) \times \prod_{i=1}^n {}^{N_i}B_{\varepsilon_i}(x_i)$. The topology on $J^n[\Sigma]$ is generated by the base $\mathfrak{B}^n = \bigcup_{\alpha\alpha'} \mathfrak{A}_{\alpha\alpha'}^n$, where $\mathfrak{A}_{\alpha\alpha'}^n = \{(\pi_n^0)^{-1}(U \times U') \mid U \times U' \subset U_\alpha \times U_{\alpha'} \text{ open}\}$.

⁸For $j^1\phi(p) \in U_{\alpha\alpha'}^1 \Rightarrow \pi_1^0(j^1\phi(p)) = (p, \phi(p)) \in U_\alpha \times U_{\alpha'} \Rightarrow j^2\phi(p) \in U_{\alpha\alpha'}^2$, and so, $U_{\alpha\alpha'}^1 \subseteq \pi_2^1(U_{\alpha\alpha'}^2)$. Conversely, $j^1\phi'(p') \in \pi_2^1(U_{\alpha\alpha'}^2) \Rightarrow j^2\phi'(p') \in U_{\alpha\alpha'}^2 \Rightarrow (p', \phi'(p')) \in U_\alpha \times U_{\alpha'} \Rightarrow$ the converse set-inequality, as required.

It follows from the construction that $\{J^n[\Sigma], \pi_n^{n-1}, \mathbb{N}\}$ forms an inverse sequence. Let $J^\infty[\Sigma] \stackrel{\text{def}}{=} \varprojlim J^n[\Sigma]$ denote the limit of the inverse sequence. Then, $\mathfrak{B}^\infty = \{(\pi^n)^{-1}(U) \mid U \in \mathfrak{B}^n \ \forall n\}$ defines a base of $J^\infty[\Sigma]$, where $\pi^n \stackrel{\text{def}}{=} p^n|J^\infty[\Sigma]$ and $p^n : \prod_{i \in \mathbb{N}} J^i[\Sigma] \rightarrow J^n[\Sigma]$ is the n th projection. Observe from [1, p. 98, proposition 2.5.1] that $J^\infty[\Sigma]$ is closed in the Cartesian product $\prod_{i \in \mathbb{N}} J^i[\Sigma]$.

The *compact* (or *weak*) C^∞ -topology on $C^\infty(\Sigma, \Sigma)$ is the topology induced by the map $j^\infty : C^\infty(\Sigma, \Sigma) \rightarrow C(\Sigma, J^\infty[\Sigma])$ defined by $f \mapsto j^\infty f \stackrel{\text{def}}{=} [f, \cdot]_\infty$ such that it is a topological imbedding. Let $\text{Diff}(\Sigma) \subset C^\infty(\Sigma, \Sigma)$ denote the set of C^∞ -diffeomorphisms on Σ . The composition mapping $\circ : \text{Diff}(\Sigma) \times \text{Diff}(\Sigma) \rightarrow \text{Diff}(\Sigma)$ given by $(f, g) \mapsto f \circ g$ defines a group structure on $\text{Diff}(\Sigma)$. Indeed, the group structure is compatible with the compact C^∞ -topology on $\text{Diff}(\Sigma)$ [3, p. 64, ex. 9]. Lastly, observe from [3, p. 38, theorem 1.6] that $\text{Diff}(\Sigma)$ is open in $C^\infty(\Sigma, \Sigma)$ (as Σ is compact implies that the weak and strong C^∞ -topology coincide).

This appendix will conclude with a brief sketch of the compact C^∞ -topology on the space Γ_2^+ of (admissible) Riemannian metrics on Σ . Let $p_\Sigma : S_2^+ \Sigma \rightarrow \Sigma$ be the symmetric covariant 2-tensor bundle over Σ and $p_{\Sigma n} : J^n[p_\Sigma] \rightarrow \Sigma$ be the C^n -jet bundle of the cross-sections of $S_2^+ \Sigma$. Then, defining $\pi_{\Sigma 1}^0 : J^1[p_\Sigma] \rightarrow \Sigma \times S_2^+ \Sigma$ as above by $j^1 q(x) \mapsto (x, q(x))$ and $\pi_{\Sigma n}^m : J^n[p_\Sigma] \rightarrow J^m[p_\Sigma]$ by $j^n q(x) \mapsto j^m q(x)$ whenever $m \leq n$, one again obtains an inverse sequence $\{J^n[p_\Sigma], \pi_{\Sigma n}^{n-1}, \mathbb{N}\}$, where $J^0[p_\Sigma] \stackrel{\text{def}}{=} \Sigma \times S_2^+ \Sigma$. Finally, let $J^\infty[p_\Sigma]$ denote the inverse limit of the sequence and set $\pi_\Sigma^n \stackrel{\text{def}}{=} p_\Sigma^n|J^\infty[p_\Sigma]$, where $p_\Sigma^n : \prod_{i \in \mathbb{N}} J^i[p_\Sigma] \rightarrow J^n[p_\Sigma]$ is the n th projection. The topology of Γ_2^+ is then defined by the (topological) imbedding $j^\infty : \Gamma_2^+ \hookrightarrow C(\Sigma, J^\infty[p_\Sigma])$.

REFERENCES

1. Engelking, R., *General Topology*, SSPM 6, Heldermann Verlag-Berlin, 1989.
2. Fischer, A. E., *Theory of Superspace*, Relativity, ed. Carmeli, Fickler, Witten.
3. Hirsch, M., *Differential Topology*, GTM 33, Springer-Verlag, Berlin, 1976.
4. Kelley, J., *General Topology*, Van Nostrand, New York, 1955.
5. Michor, P., *Manifolds of Differentiable Mappings*, vol. 8, Shiva Maths Series, Kent, 1980.
6. Rovelli, C., *Class. Quantum Grav* **8** (1991), 1613–1675.
7. Rovelli, C. and Smolin, L., *Nuc. Phys* **B331** (1990), 80–152.
8. Saunders, D., *The Geometry of Jet Bundles*, LMS 142, Cambridge University Press, 1989.